

An equivalence relation of boundary/initial conditions, and the infinite limit properties.

KAZUHIKO MINAMI

*Graduate School of Mathematics, Nagoya University,
Nagoya, 464-8602, JAPAN*

February 2, 2008

The 'n-equivalences' of boundary conditions of lattice models are introduced and it is derived that the models with n-equivalent boundary conditions result in the identical free energy. It is shown that the free energy of the six-vertex model is classified through the density of left/down arrows on the boundary. The free energy becomes identical to that obtained by Lieb and Sutherland with the periodic boundary condition, if the density of the arrows is equal to $1/2$. The relation to the structure of the transfer matrix and a relation to stochastic processes are noted.

minami@math.nagoya-u.ac.jp

Two configurations $\{s_i\}$ and $\{s'_i\}$ where the corresponding local variables are the same: $s_i = s'_i$, except a finite number of them can be regarded as being identical when we consider the thermodynamic limit properties. This is also true when the differences approach infinity but remain of the lower orders than bulk quantities such as the total number of sites or the total energy contributions. It follows that boundary conditions, for example, of lattice models are usually irrelevant to the thermodynamics.

On the other hand, a six-vertex model with a specific boundary condition was solved [1] and it is found that the free energy of it is not identical to the free energy previously obtained [2, 3, 4] with the cyclic boundary condition. This difference of two free energies comes from the fact that the configurations of the six-vertex model should satisfy the six-vertex restriction, and therefore the boundary condition is still relevant in the thermodynamic limit, even if the energy contributions coming from the boundary sites are irrelevant in the limit.

In this short note an equivalence relation of boundary conditions is introduced. It is generally derived that two boundary conditions yield the same free energy if they belong to the same equivalence class, even in models with long-ranged restrictions. Several related properties will also be discussed.

First let us define the equivalence relation. Let us consider classical lattice models where each local variables takes one of a finite number of states. We assume that the number of boundary sites is of a lower order than the total number of sites. Let us consider the set of sites which lie in the lattice and can be reached from the boundary sites by just n steps (n bonds) at minimum; we call these sites the n -boundary sites. Let us consider the set of bonds between $(n-1)$ - and n -boundary sites, and call them the n -boundary bonds. At last let us consider the n -boundary sites together with the n -boundary bonds and call them the n -boundary, and also call the configurations on the n -boundary the n -boundary configurations. Let $\{\Gamma_i\}$ be the set of all the possible configurations on the n -boundary with a boundary condition Γ on the actual boundary of the lattice. Two boundary conditions Γ and Γ' are called n -equivalent when $\{\Gamma_i\} = \{\Gamma'_i\}$ as a set of n -boundary configurations.

Then we can show the following fact. Suppose that the boundary conditions Γ and Γ' are n -equivalent throughout the thermodynamic limit with some finite n . Then the two free energies with Γ and Γ' are identical. Proof: Let us write the partition function as $Z = \sum_i B_i Z_i$. The factor Z_i is the partition function from the set of local variables on the lattice inside the n -boundary with fixed n -boundary configuration Γ_i . The factor B_i is the

partition function from the other local variables with Γ and Γ_i . Then the factor $B_i Z_i$ is the partition function of the system with the Γ and fixed Γ_i . From the assumption that n is always finite, we obtain $\log Z_i = -\beta N f_i + o(N)$ and $\log B_i = o(N)$ where N is the total number of sites. The index i runs from 1 to $i_{\max}$ where $i_{\max}$ is the number of the possible configurations on the n -boundary with the boundary condition Γ . We have $i_{\max} \leq O(r^{N'})$ where r is a constant and N' is the number of sites on the n -boundary. Then we obtain

$$\begin{aligned} \frac{1}{N} \log Z &= \frac{1}{N} \log \left(\sum_i B_i Z_i \right) \\ &= \frac{1}{N} \log B_1 Z_1 \left[1 + \sum_{i=2}^{m_1} \frac{B_i}{B_1} \frac{Z_i}{Z_1} + \sum_{i=m_1+1}^{i_{\max}} e^{-\beta N(f_i - f_1) + o(N)} \right] \\ &\rightarrow -\beta f_1 \quad (N \rightarrow \infty), \end{aligned}$$

where $f_i = f_1$ ($1 \leq i \leq m_1$), $f_1 < f_i$ ($m_1 + 1 \leq i$). ■

All the boundary conditions are 1-equivalent to each other for the Ising models because the spin states are independent of their nearest-neighbors. This results in the following trivial fact that the free energies of the Ising models with finite range interactions are independent of the boundary conditions.

This equivalence can be used in a slightly generalized form. We can concentrate on a part of the boundary (*e.g.* one of the four edges of the rectangle, this will be considered below) with a condition Γ , and introduce the corresponding n -boundary and the corresponding n -equivalence.

Let us consider the six-vertex model on a rectangular lattice with h rows and w columns. We assign an arrow on each bond. Two arrows in and the other two out at each site. Then there exist six types of possible local arrow arrangements as shown in Fig.1, and we call them the six vertices. Each vertex has finite energy, the energy is assumed to be unchanged reversing all the four arrows around the site. One can assign a line to a bond if the arrow on the bond points down or points left, as shown again in Fig.1, and there exists a one-to-one correspondence between configurations of arrows and configurations of continuous lines on the lattice. The possible six arrow arrangements satisfy the restriction that the lines do not intersect each other and continue beginning from one bond on the boundary of the lattice to another bond on the boundary. The total number of lines is hence conserved when the boundary configuration is fixed.

Let us assume that the line configuration on the first row of vertical bonds (upper edge of the rectangle) is fixed and identical to that on the last row, and also assume that the line configuration on the first column of horizontal bonds (the right edge of the rectangle) is fixed and identical to that on the last column. In this case the number of lines is conserved in each step moving from the first row to the last row, and also from the first column to the last column. Let us assume that there are always n_1 (n_2) lines on the first p_0 boundary bonds on the upper (right) edge and n_1 (n_2) lines on the next p_0 bonds on the edges, and vice versa. In this case we introduce the boundary line density ρ_i as $\rho_i = n_i/p_0$ where $i = 1$ and 2 . Fix a sequence of finite rectangles approaching to the thermodynamic limit. Then the free energy of the six-vertex model is determined by ρ_1 and ρ_2 : $f = f(\rho_1, \rho_2)$. Proof: If $\rho_2 = 0$ the free energy is equal to $(1 - \rho_1)\epsilon_1 + \rho_1\epsilon_2$ (and similarly if $\rho_1 = 0$). As for the cases with the densities equal to one, we can use the fact that the free energy with $(\rho_1, \rho_2) = (1 - \rho, 1 - \rho')$ is identical to that with $(\rho_1, \rho_2) = (\rho, \rho')$, which comes from the symmetry of the vertex energies. If both of ρ_1 and ρ_2 satisfy $0 < \rho_i < 1$, consider an edge with the density ρ_1 and the corresponding n -boundary. It is easy to convince that all the boundary line configurations on the edge with the density ρ_1 are n -equivalent to each other, where $n = kp_0$ is now a fixed constant proportional to p_0 . This argument is valid respectively for the other three edges. ■

It can be derived [5], with the use of the n -equivalence, in the case of the six-vertex model with more general situation, that $f(1/2, 1/2) = f_{\text{LS}}$ where f_{LS} is the free energy obtained [2, 3, 4] with the cyclic boundary condition in both the horizontal and the vertical directions.

This argument can be applied to line conserved models on other lattices. The argument is also easily generalized to models with l_i lines on each bond i , where $l_i = 0, 1, \dots, q - 1$, that is the q state vertex models. On the square lattice, for example, with $q = 3$ we have the 19-vertex model.

A kind of SOS model is equivalent [6] to a q state vertex model and also equivalent to a class of spin S classical and quantum spin chains, where the total magnetization corresponds to the number of lines.

Next we note the relation between the n -equivalence and the structure of the transfer matrix. Let us consider the six-vertex model on the rectangle and assume the cyclic boundary condition in the horizontal direction, i.e. the six-vertex model on a cylinder. The free energy is obtained from the maximum eigenvalue of the transfer matrix V . All the matrix elements of V are non-negative because each element is a sum of Boltzmann weights. The

matrix V is block-diagonalized according to the number of lines m which is conserved from row to the next row, and let V_m be the corresponding block element. It is easy to convince that all the line configurations with m lines, $m \leq w$, on the upper edge are n -equivalent to each other, where $n = kw$ and k is a positive integer. This means that there are allowed line configurations on the lattice with arbitrary fixed boundary line configurations with m lines on the first row and other arbitrary line configurations with m lines on the last row, if h is larger than $2kw$. Then all the matrix elements of V_m^h are positive, and hence V_m^h is irreducible, because any matrix is irreducible if all of its elements are positive. We can thus conclude that V_m is irreducible, because V_m^h cannot be irreducible if V_m is not. This fact means that the transfer matrix is irreducible if the bases (line configurations on the row) used for the matrix representation are n -equivalent to each other. We can also say that all the line configurations on the first row with m lines result in the same free energy in the limit $h \rightarrow \infty$ with fixed w , because they are kw -equivalent and kw is finite. The free energies remains of course identical when we take the limit $w \rightarrow \infty$ afterwards.

At last we will make a comment on the relation with stochastic processes. In this case, V is a stochastic matrix where the sum of the elements of each row is unity that is the total sum of probabilities. It is known that the maximum eigenvalues of such matrix are always unity and the absolute values of the other eigenvalues do not exceed unity. Let us consider a stochastic system characterized by a finite number of variables $(l_1, \dots, l_s)$, with a kind of conservation law $\sum l_i = \text{const}$. This kind of stochastic processes correspond to the line-conserved lattice models. The stochastic matrix is block diagonalized for each set of n -equivalent initial conditions. Each block element is irreducible and the limiting state is again classified according to the n -equivalence.

References

- [1] V.Korepin and P.Zinn-Justin: J.Phys.A **33** (2000) 7053
- [2] E.H.Lieb: Phys.Rev.Lett.**18** (1967) 1046, **19** (1967) 108
- [3] E.H.Lieb: Phys.Rev. **162** (1967) 162
- [4] B.Sutherland: Phys.Rev.Lett.**19** (1967) 103
- [5] K.Minami: in preparation
- [6] K.Sogo, Y.Akutsu, T.Abe: Prog.Theor.Phys.**70** (1983) 739




Fig.1 Six vertices, corresponding lines and energies.